ENFERMEDAD RENAL CRÓNICA

GUÍAS KDIGO 2024

Mar Castiella Bañales
MIR 4º año

ÍNDICE

DETECCIÓN

CLASIFICACIÓN

EVALUAR EL RIESGO

TRATAMIENTO

MANEJO COMPLICACIONES

REMITIR A NEFROLOGÍA

DEFINICIÓN

La ERC se define como el daño estructural o funcional del riñón mantenido más de 3 meses:

MARCADORES DAÑO RENAL

- Albuminuria (albumina/creatinina > 30 mg/g [>3 mg/mmol])
- Alteraciones en el sedimento urinario
- Hematuria
- Alteraciones tubulares con anomalías electrolíticas
- Alteraciones histológicas
- Alteraciones estructurales detectadas con imagen (poliquistosis renal..)
- Antecedentes de trasplante renal

DISMINUCIÓN FGE

FGE < 60 mL/min/1.73m²

Prevalencia global de ERC 9,1%
¡Conciencia de enfermedad < 10 %!

DETECCIÓN ERC

CRIBADO A PERSONAS DE RIESGO

Medir:

1. Tas
2. Coc
3. Otr

Revisión de **mediciones previas alteradas**

Hallazgos de imagen como la reducción del tamaño renal o el grosor cortical

Hallazgos histológicos de fibrosis y atrofia

Condiciones clínicas que lo sugieran (HTA o DM de larga evolución mal controlados)

Hipertensión arterial
Diabetes
Enfermedad CV
Obesidad
Antecedentes de IRA
Antecedentes familiares
Comorbilidades
(nefrotóxicos, lupus...)

FGE < 60 mL/min/1.73 m² +/- CAC > 30 mg/g +/- presencia de otros marcadores de daño renal **durante 3 meses**
evidencia de cronicidad

Clasificación
Establecer la causa
Calcular riesgo de progresión
Iniciar tratamiento

FGE < 30 mL/min/1.73 m² +/- CAC > 30 mg/g agudo

Daño renal agudo

CLASIFICACIÓN

Sistema CGA:

- Causa
- FGE
- Albuminuria

Esta clasificación determina la **severidad y el riesgo** de:

**PROGRESIÓN
EVENTOS CV
MORTALIDAD**

CAUSAS

Table 6 | Guidance for the selection of additional tests for evaluation of cause

Test category	Examples	Comment or key references
Imaging	Ultrasound, intravenous urography, CT kidneys ureters bladder, nuclear medicine studies, MRI	Assess kidney structure (i.e., kidney shape, size, symmetry, and evidence of obstruction) for cystic disease and reflux disease. Evolving role of additional technologies (e.g., 3D ultrasound)
Kidney biopsy	Ultrasound-guided percutaneous	Usually examined by light microscopy, immunofluorescence, and electron microscopy, and, in some situations, may include molecular diagnostics. Used for exact diagnosis, planning treatment, assessing activity and chronicity of disease, and likelihood of treatment response; may also be used to assess genetic disease
Laboratory tests: serologic, urine tests	Chemistry including acid-base and electrolytes, serologic tests such as anti-PLA2R, ANCA, anti-GBM antibodies Serum-free light chains, serum, and urine protein electrophoresis/immunofixation Urinalysis and urine sediment examination	Refer to <i>KDIGO 2021 Clinical Practice Guideline for the Management of Glomerular Diseases</i> ²² Increasing recognition of the role of light chains in kidney disease even in the absence of multiple myeloma (monoclonal gammopathy of renal significance [MGRS]) ⁹⁸ Presence of persistent hematuria or albuminuria is critical in determining differential diagnosis
Genetic testing	<i>APOL1, COL4A3, COL4A4, COL4A5, NPHS1, UMOD, HNF1B, PKD1, PKD2</i>	Evolving as a tool for diagnosis, increased utilization is expected. Recognition that genetic causes are more common and may present without classic family history ^{99,100}

ANCA, antineutrophil cytoplasmic antibody; *APOL1*, apolipoprotein 1; *COL4A*, type IV collagen alpha chain; CT, computed tomography; GBM, glomerular basement membrane; *HNF1B*, hepatocyte nuclear factor 1B; MRI, magnetic resonance imaging; *NPHS1*, congenital nephrotic syndrome; *PKD1*, polycystic kidney disease-1; *PKD2*, polycystic kidney disease-2; PLA2R, M-type phospholipase A2 receptor; *UMOD*, uromodulin.

Recommendation 1.1.4.1: We suggest performing a kidney biopsy as an acceptable, safe, diagnostic test to evaluate cause and guide treatment decisions when clinically appropriate (2D).

CAUSAS

50%

ENFERMEDAD
GLOMERULAR

Diabetes, enfermedades autoinmunes,
medicaciones, neoplasias (incluye
amiloidosis),...

ENFERMEDAD
TUBULOINTERSTICIAL

Infecciones sistémicas, enfermedades
autoinmunes, sarcoidosis, medicamentos,
neoplasias (mieloma),...

ENFERMEDAD
VASCULAR

Ateroesclerosis, hipertensión,
isquemia, vasculitis,...

ENFERMEDAD QUÍSTICA
CONGENITA

Enfermedad renal poliquística,
enfermedad de Fabry, síndrome de
Alport...

EVALUACIÓN DE TFG

GFR category	GFR (ml/min per 1.73 m ²)	Terms
G1	≥90	Normal or high
G2	60-89	Mildly decreased*
G3a	45-59	Mildly to moderately decreased
G3b	30-44	Moderately to severely decreased
G4	15-29	Severely decreased
G5	<15	Kidney failure

FG ≤ 60 ML/MIN/1.73 M2

CAÍDA FGE DE 0.6 - 0.8 ML AÑO

EVALUACIÓN DE ALBUMINURIA

Category	AER (mg/24 hours)	ACR (approx. equivalent)		Terms
		(mg/mmol)	(mg/g)	
A1	<30	<3	<30	Normal to mildly increased
A2	30-300	3-30	30-300	Moderately increased*
A3	>300	>30	>300	Severely increased†

EVALUAR EL RIESGO

KDIGO: Prognosis of CKD by GFR and albuminuria categories

				Persistent albuminuria categories		
				Description and range		
				A1	A2	A3
				Normal to mildly increased	Moderately increased	Severely increased
				<30 mg/g <3 mg/mmol	30–300 mg/g 3–30 mg/mmol	>300 mg/g >30 mg/mmol
GFR categories (ml/min/1.73 m ²) Description and range	G1	Normal or high	≥90	Green	Yellow	Orange
	G2	Mildly decreased	60–89	Green	Yellow	Orange
	G3a	Mildly to moderately decreased	45–59	Yellow	Orange	Red
	G3b	Moderately to severely decreased	30–44	Orange	Red	Red
	G4	Severely decreased	15–29	Red	Red	Red
	G5	Kidney failure	<15	Red	Red	Red

Green: low risk (if no other markers of kidney disease, no CKD); Yellow: moderately increased risk; Orange: high risk; Red: very high risk. GFR, glomerular filtration rate.

- **PROGRESIÓN**
- **EVENTOS CV**
- **MORTALIDAD**

EVALUAR EL RIESGO

- El riesgo se aproxima con el mapa pero no es absoluto.

- En ries

Age (Yrs)

GFR (ML/Min/1.73M2)

Urine Albumin: Creatinine Ratio Units

?

?

Select

NEXT

de

ía IgA)...

logo.
disciplinario.
para trasplante y

RETRASO DE LA PROGRESIÓN

MANEJO NO FARMACOLÓGICO

CESE TABÁQUICO

DIETA

Dieta variada y saludable
Aumentar consumo de
alimentos de origen vegetal
Evitar ultraprocesados

EJERCICIO FÍSICO

150 minutos a la semana

Ingesta de proteínas de 0,8
g/kg/día

*No restringir en niños, pacientes frágiles,
sarcopénicos..

Ingesta de sodio <2g (<5g sal)

MANEJO FARMACOLÓGICO

Inhibidores del sistema renina-angiotensina (IECA ó ARA II)

FGE > 15

- Todo paciente con albuminuria A3 (1B)
- Todo paciente con albuminuria A2 + diabetes (1B)
 - Considerar en albuminuria A2 sin diabetes (2C)

Considerar en albuminuria A1 para indicaciones específicas (HTA, IC FEVI disminuida)

- Utilizar la dosis más alta tolerada
- Control de TA, creatinina sérica y potasio en 2-4 semanas

MANEJO FARMACOLÓGICO

Inhibidores del cotransportador sodio-glucosa-2 (SGLT2i)

FGE > 20

- Todo paciente con ERC + diabetes (1A)
- Todo paciente con ERC + IC (1A)
- Todo paciente con CAC > 200mg/g (1A)
 - Considerar en FGE 20-45 sin albuminuria (2B)

- Continuar si FGe descende < 20
- Suspender en situaciones de riesgo de cetosis (ayuno, cirugía..)

MANEJO FARMACOLÓGICO

Antagonistas de los receptores de mineralocorticoides (ARM)

Finerenona

FGE > 25

- **Todo paciente con ERC + diabetes + albuminuria A2,A3 + K normal** a pesar de dosis máxima tolerada de IECA/ARAII (2A)

- **¡Vigilar K!**
 - **Suspender si K > 5,5 mmol/l**
- **Se pueden combinar con SGLT2i**

MANEJO DE COMPLICACIONES

ACIDOSIS METABÓLICA

PREVALENCIA

G3: 7,7 %
G5: 38,3%

BICARBONATO

- No hay clara evidencia de su beneficio
- Considerar si $\text{HCO}_3^- < 18 \text{ mmol/l}$

HIPERPOTASEMIA

PREVALENCIA

G3: 5-9%.

G5: 24-34%

1st line: Address correctable factors	<ul style="list-style-type: none">• Review non-RASi <u>medications</u> (e.g. NSAIDs, trimethoprim)• Assess <u>dietary</u> potassium intake (dietary referral) and consider appropriate moderation of dietary potassium intake
2nd line: Medications	Consider: <ul style="list-style-type: none">• Appropriate use of diuretics• Optimize serum bicarbonate levels• Licensed potassium exchange agents
3rd line: Last resort	<ul style="list-style-type: none">• Reduce dose or discontinue RASi/MRA (Discontinuation is associated with increased cardiovascular events. Review and restart RASi or MRA at a later date if patient condition allows.)

Figure 32 | Actions to manage hyperkalemia (potassium >5.5 mmol/l) in chronic kidney disease. MRA, mineralocorticoid receptor antagonists; NSAID, nonsteroidal anti-inflammatory drug; RASi, renin-angiotensin system inhibitors.

HIPERPOTASEMIA

Table 26 | Medications associated with increased risk of hyperkalemia

Class	Mechanism	Example
ACEi	Inhibit conversion of angiotensin I to angiotensin II	Captopril, lisinopril, perindopril, etc.
ARB	Inhibit activation of angiotensin I receptor by angiotensin II	Losartan, irbesartan, candesartan, etc.
Aldosterone antagonist	Block aldosterone receptor activation	Spirolactone, eplerenone, and finerenone
β -Adrenergic receptor blocker	Inhibit renin release	Propranolol, metoprolol, and atenolol
Digitalis glycoside	Inhibit Na^+ - K^+ -ATPase, necessary for collecting duct K^+ secretion	Digoxin
Heparin	Reduced production of aldosterone	Heparin sodium
Potassium-sparing diuretic	Block collecting duct apical Na^+ channel, decreasing gradient for K^+ secretion	Amiloride and triamterene
NSAIDs	Inhibit synthesis of prostaglandin E and prostacyclin, inhibiting renin release	Ibuprofen, naproxen, diclofenac, etc.
CNI	Inhibit Na^+ - K^+ -ATPase, necessary for collecting duct K^+ secretion	Cyclosporine and tacrolimus
ns-MRA	Block MR-mediated Na^+ reabsorption	Finerenone
Other	Block collecting duct apical Na^+ channel, decreasing gradient for K^+ secretion	Trimethoprim and pentamidine

ACEi, angiotensin-converting enzyme inhibitor; ARB, angiotensin II receptor blocker; ATP, adenosine triphosphate; CNI, calcineurin inhibitor; K^+ , potassium; Na^+ , sodium; NSAID, nonsteroidal anti-inflammatory drug; ns-MRA, nonsteroidal mineralocorticoid receptor antagonist.

Data from Weiner *et al.*⁵⁷⁶ and Kidney Disease: Improving Global Outcomes Diabetes Work Group.²³

HIPERPOTASEMIA

Figure 33 | Potassium absorption rates of plant-based, animal-based, and processed foods. Data from Picard K, Griffiths M, Mager DR, Richard C. Handouts for low-potassium diets disproportionately restrict fruits and vegetables. *J Ren Nutr.* 2021;31:210–214.⁵⁹²

ANEMIA

PREVALENCIA

G3: 14,9 %

G5: 60,7 %

- **No** atribuir a ERC. Hacer un **adecuado abordaje diagnóstico**.
- Si finalmente es anemia renal, valorar Eritropoyetina (si Hb<10 g/dL)

ENFERMEDAD MINERAL-ÓSEA

- Alteraciones en fósforo, calcio, PTH, vitamina D y FGF23.
- Condiciona múltiples enfermedades óseas.

HIPERURICEMIA

- Hiperuricemia sintomática (tras primer episodio de gota): tratamiento preferiblemente con Alopurinol.
- Evitar AINES; mejor Colchicina o corticoides.

ENFERMEDAD CARDIOVASCULAR

- La **enfermedad c**
- los pacientes con
- Objetivo de PAS <

- **Pacientes con ERC**
- **ERC sin diabetes o EA**

KDIGO: Prognosis of CKD by GFR and albuminuria categories

				Persistent albuminuria categories		
				Description and range		
				A1	A2	A3
				Normal to mildly increased	Moderately increased	Severely increased
				<30 mg/g <3 mg/mmol	30–300 mg/g 3–30 mg/mmol	>300 mg/g >30 mg/mmol
GFR categories (ml/min/1.73 m ²) Description and range	G1	Normal or high	≥90			
	G2	Mildly decreased	60–89			
	G3a	Mildly to moderately decreased	45–59			
	G3b	Moderately to severely decreased	30–44			
	G4	Severely decreased	15–29			
	G5	Kidney failure	<15			

Green: low risk (if no other markers of kidney disease, no CKD); Yellow: moderately increased risk; Orange: high risk; Red: very high risk. GFR, glomerular filtration rate.

mortalidad en

il/ 10 a/	<p>Riesgo alto</p> <p>LDL < 70</p>
	<p>Riesgo muy alto</p> <p>LDL < 55</p>

5):429.e1–429.e104

FIBRILACIÓN AURICULAR

- La prevalencia de FA es mayor en ERC. Las complicaciones son mayores (ACV).
- Se recomiendan los **anticoagulantes orales no antagonistas de la vitamina K (ACODs)** frente a los antagonistas de la vitamina K (warfarina).
 - *Ajustar dosis por función renal.*
- En diálisis no hay clara evidencia del beneficio de la anticoagulación.

GESTIÓN DE LA MEDICACIÓN Y ADMINISTRACIÓN DE FÁRMACOS

Table 31 | Key examples of common medications with documented nephrotoxicity and, where available, selected non-nephrotoxic alternatives

Nephrotoxic medication	Potential non-nephrotoxic alternatives
<p>Analgesics NSAIDs: nephrotoxic effects include a decrease in GFR through a reduction in prostaglandin-dependent kidney blood flow, allergic interstitial nephritis (AIN), and nephrotic syndrome⁷²⁵</p>	Acetaminophen
<p>Antimicrobials Aminoglycosides: accumulates in the proximal tubular cells and disrupts phospholipid metabolism, resulting in cell apoptosis and acute tubular necrosis (ATN)^{731,732}</p>	Cephalosporins and carbapenems
<p>Vancomycin: unclear cause of nephrotoxicity, but likely related to ATN and possible AIN^{731,732}</p>	Linezolid and daptomycin ⁷³¹
<p>Sulfamethoxazole-trimethoprim: AIN, ATN, crystalluria within the distal convoluted tubule and reversible inhibition of tubular creatinine secretion⁷³¹</p>	Clindamycin + primaquine, pentamidine, and atovaquone
<p>Gastrointestinal medications Proton pump inhibitors: may result in AKI and CKD due to tubulointerstitial nephritis and AIN^{733,734}</p>	H2-receptor antagonists Ranitidina, Cimetidina
<p>Cardiovascular medications Warfarin: glomerular hemorrhage, oxidative stress causing kidney tubular damage, and direct effects on kidney vascular calcification by vitamin K-dependent alterations of matrix Gla protein^{735,736}</p>	Non-vitamin K antagonist oral anticoagulants
<p>Other Lithium: nephrogenic diabetes insipidus as well as CKD from chronic tubulointerstitial nephropathy⁷³⁷</p>	Aripiprazole, lamotrigine, quetiapine, valproate

CKD, chronic kidney disease; GFR, glomerular filtration rate; NSAID, nonsteroidal anti-inflammatory drug.

43 %

REMITIR A NEFROLOGÍA

CONCLUSIONES

- La ERC es una patología muy prevalente pero silente.
- Es posible tener ERC con una tasa de filtración glomerular normal.
- Importante estimar el riesgo de progresión, eventos CV y mortalidad.
- Tratamiento con IECAS si albuminuria.
- Tratamiento con SGLT2i si diabetes, albuminuria o $FG < 45$.
- Finerenona como tratamiento novedoso cuando no se consiguen objetivos con IECAS.

BIBLIOGRAFÍA

- KDIGO 2024: Clinical Practice Guideline for the Management of Chronic Kidney Disease
- 2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk

**iMUCHAS
GRACIAS!**